CORRECTION Open Access

Correction to: Cross-sectional study of associations between normal body weight with central obesity and hyperuricemia in Japan



Takako Shirasawa^{1*}, Hirotaka Ochiai¹, Takahiko Yoshimoto¹, Satsue Nagahama², Akihiro Watanabe¹, Reika Yoshida¹ and Akatsuki Kokaze¹

Correction to: BMC Endocr Disord (2020) 20:2 https://doi.org/10.1186/s12902-019-0481-1

After publication of this article [1], it was brought to our attention that there is an error in the Table 5, which "Cases" should be revised to "HU" in Table 5. The original publication has been corrected.

Author details

¹Department of Hygiene, Public Health and Preventive Medicine, Showa University School of Medicine, 1-5-8 Hatanodai, Shinagawa-ku, Tokyo 142-8555, Japan. ²All Japan Labor Welfare Foundation, 6-16-11 Hatanodai, Shinagawa-ku, Tokyo 142-0064, Japan.

Published online: 21 February 2020

Reference

 Shirasawa, et al. BMC Endocr Disord. 2020;20:2. https://doi.org/10.1186/ s12902-019-0481-1.

The original article can be found online at https://doi.org/10.1186/s12902-019-0481-1

Full list of author information is available at the end of the article



^{*} Correspondence: shirasawa@med.showa-u.ac.jp

¹Department of Hygiene, Public Health and Preventive Medicine, Showa University School of Medicine, 1-5-8 Hatanodai, Shinagawa-ku, Tokyo 142-8555, Japan