CORRECTION Open Access

Correction: Inadequate care and excessive overprotection during childhood are associated with the presence of Diabetes Mellitus in adulthood in a general Japanese population: a cross-sectional analysis from the Hisayama Study

Mao Shibata^{1,2,3*}, Masako Hosoi^{1,4}, Kozo Anno¹, Naoki Hirabayashi^{2,4}, Yoichiro Hirakawa^{2,5}, Hiroshi Kawata⁴, Rie Iwaki¹, Rvoko Sawamoto⁴, Nobuvuki Sudo^{1,4} and Toshiharu Ninomiva^{2,3}

Correction: BMC Endocr Disord 23, 222 (2023) https://doi.org/10.1186/s12902-023-01474-4

Following publication of the original article [1], the authors reported an error in the number of participants in Fig. 2.

The original article can be found online at https://doi.org/10.1186/s12902-023-01474-4.

*Correspondence:

Mao Shibata

shibata.mao.276@m.kyushu-u.ac.jp

⁵ Department of Medicine and Clinical Science, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.


¹ Department of Psychosomatic Medicine, Kyushu University Hospital, Fukuoka, Japan

² Department of Epidemiology and Public Health, Graduate School of Medical Sciences, Kyushu University, Fukuoka 812-8582, Japan


³ Division of Research Management, Center for Cohort Studies, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan

⁴ Department of Psychosomatic Medicine, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan

The original Fig. 2 was:



The updated Fig. 2 should read:



The original article [1] has been updated.

Published online: 21 November 2023

Reference

 Shibata M, Hosoi M, Anno K, et al. Inadequate care and excessive overprotection during childhood are associated with the presence of diabetes mellitus in adulthood in a general Japanese population: a cross-sectional analysis from the Hisayama Study. BMC Endocr Disord. 2023;23:222. https://doi.org/10.1186/s12902-023-01474-4.